



## inSite Digestive Healthcare COLONOSCOPY PREPARATION SHEET: Clenpiq

Facility:  Check with your gastro office if you're not CERTAIN WRITE IT DOWN

Procedure Date: \_\_\_\_\_ Check In Time: \_\_\_\_\_

5 Days Prior	4 Days Prior	3 Days Prior	2 Days Prior	One Day Prior	Colonoscopy Day
<ul style="list-style-type: none"> <li><input type="checkbox"/> Arrange for a ride to and from Facility</li> <li><input type="checkbox"/> Go to pharmacy &amp; purchase Clenpiq by prescription, use coupon if available</li> <li><input type="checkbox"/> For sore bottom:               <ul style="list-style-type: none"> <li>--plain or aloe baby wipes</li> <li>--Desitin or A&amp;D ointment</li> </ul> </li> <li><input type="checkbox"/> Stop taking Coumadin or Plavix. Follow the instructions of your physician if alternative blood thinners are needed. Do not proceed without talking to your Gastro doctor if taking <b>Brilinta or Effient</b></li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Last chance to cancel the procedure without a cancellation fee</li> <li><input type="checkbox"/> For Co-payment questions call your Gastro doctor's office</li> <li><input type="checkbox"/> Stop supplements, vitamins, probiotics, iron</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Review diet instructions for the coming 3 days. Plan meals and shop to comply with instructions</li> <li><input type="checkbox"/> Confirm your ride to and from facility</li> <li><input type="checkbox"/> Stop any fiber supplements</li> <li><input type="checkbox"/> Eat well balanced meals <b>BUT AVOID:</b> <ul style="list-style-type: none"> <li>- Raw fruits</li> <li>- Raw vegetables</li> <li>- Nuts or popcorn</li> <li>- Foods with small seeds such as tomato, sesame or melon seeds</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Drink AT LEAST 8 glasses of water or clear liquid during the day</li> <li><input type="checkbox"/> Eat well balanced meals <b>BUT AVOID:</b> <ul style="list-style-type: none"> <li>- Raw fruits</li> <li>- Raw vegetables</li> <li>- Nuts or popcorn</li> <li>- Foods with small seeds such as tomato, sesame or melon seeds</li> </ul> </li> <li><input type="checkbox"/> <b>No solid food after midnight</b></li> <li><input type="checkbox"/> Stop taking Eliquis, Pradaxa, Xarelto, Savasa after the morning dose (48 hours prior to procedure).</li> <li><input type="checkbox"/> Do NOT stop low dose (81mg) aspirin taken for heart or stroke reasons</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Drink at least 8 glasses of water and other clear liquids during the day</li> <li><input type="checkbox"/> <b>Follow a strict clear diet ALL DAY</b></li> <li><input type="checkbox"/> <b>5 or 6 pm: Start drinking 1 of the bottles of Clenpiq and follow with 5 or more 8 oz cups of clear fluids (cup included in kit)</b></li> <li><input type="checkbox"/> Can have further clear fluids as you wish</li> <li><input type="checkbox"/> If you have any nausea take a 15-30 minute break then resume drinking clear fluids</li> <li><input type="checkbox"/> Verify that you have info you'll need to bring with you to facility</li> <li><input type="checkbox"/> Do not schedule any important activities on the day of procedure</li> <li><input type="checkbox"/> <b>You may not drive or go home unaccompanied in a taxi or UBER etc after your procedure</b></li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> <b>Beginning 4 or 5 hours before leaving home for your colonoscopy, start drinking 2nd bottle of Clenpiq, followed by at least 4 more 8 oz cups of clear fluids</b></li> <li><input type="checkbox"/> <b>After the last dose, do not eat or drink anything for 3 hours</b></li> <li><input type="checkbox"/> You may take regular prescriptions with small sips of water (but no blood thinners)</li> <li><input type="checkbox"/> If you take insulin shots, follow the instructions for lowered dose you were given for procedure day</li> <li><input type="checkbox"/> <b>Be sure to bring:</b> <ul style="list-style-type: none"> <li>• Insurance card or Medicare card</li> <li>• Identification (CA driver's license, or ID card)</li> <li>• Cash, check or credit card (Visa or Master Card) for co-payment and/or deductible payment. (For co-payments or deductible amount, please contact your facility)</li> </ul> </li> <li><input type="checkbox"/> Allow 2-3 hours at the facility before your driver will pick you up. Staff will call your driver to coordinate</li> <li><input type="checkbox"/> Normal diet after you are home</li> </ul>

Two days prior: Consider stopping all casual (not Dr. recommended) Aspirin or NSAIDS-no ibuprofen, Advil, Aleve, Naprosyn, Excedrin---only Tylenol is okay or meds for migraine



**inSite Digestive Healthcare**  
**COLONOSCOPY PREPARATION SHEET: Clenpiq**

Facility:  Check with your gastro office if you're not CERTAIN WRITE IT DOWN

Procedure Date: \_\_\_\_\_ Check In Time: \_\_\_\_\_

--	--	--	--	--	--

Two days prior: Consider stopping all casual (not Dr. recommended) Aspirin or NSAIDS-no ibuprofen, Advil, Aleve, Naprosyn, Excedrin---only **Tylenol is okay or meds for migraine**